

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
12 July 2001 (12.07.2001)

PCT

(10) International Publication Number
WO 01/49673 A2

(51) International Patent Classification?: C07D 311/00

11435 Belshire Lane, San Diego, CA 92126 (US). KHAM-MUNGKHUNE, Sak; 3530 42nd Street, #4, San Diego, CA 92105 (US).

(21) International Application Number: PCT/US00/35671

(22) International Filing Date:
29 December 2000 (29.12.2000)

(74) Agents: MISROCK, Leslie, S. et al.; Pennic & Edmonds LLP, 1155 Avenue of the Americas, New York, NY 10036 (US).

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
09/475,776 30 December 1999 (30.12.1999) US
09/492,939 27 January 2000 (27.01.2000) US
09/611,156 6 July 2000 (06.07.2000) US

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.



(71) Applicant: SIGNAL PHARMACEUTICALS, INC.
[US/US]; 5555 Oberlin Drive, San Diego, CA 92121 (US).

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,

(72) Inventors: BHAGWAT, Shripad, S.; 5015 Ashley Falls Court, San Diego, CA 92130 (US). MCKIE, Jeffrey, A.;

[Continued on next page]

(54) Title: COMPOUNDS AND METHODS FOR MODULATION OF ESTROGEN RECEPTORS



(57) Abstract: Compounds that modulate gene expression through the estrogen receptor (ER) are disclosed having formula (I), as well as pharmaceutical compositions containing the same wherein R_1 , R_2 , R_3 , n and p are as defined here. In a specific embodiment, the compounds are selective modulators for ER- β over ER- α . Methods are also disclosed for modulating ER- β in cells and/or tissues expressing the same, including cells and/or tissues that preferentially express ER- β . More generally, methods for treating estrogen-related conditions are also disclosed, including conditions such as breast cancer, testicular cancer, osteoporosis, endometriosis, cardiovascular disease, hypercholesterolemia, prostatic hypertrophy, prostatic carcinomas, obesity, hot flashes, skin effects, mood swings, memory loss, urinary incontinence, hairloss, cataracts, natural hormonal imbalances, and adverse reproductive effects associated with exposure to environmental chemicals.

WO 01/49673 A2



IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

Published:

— *Without international search report and to be republished upon receipt of that report.*

COMPOUNDS AND METHODS FOR MODULATION OF ESTROGEN RECEPTORS

CROSS-REFERENCE TO RELATED APPLICATION

This application claims the benefit of U.S. Provisional Application No. 60/114,472, filed December 30, 1998, and is a continuation-in-part of PCT/US99/31290 filed December 30, 1999, U.S. Application No. 09/475,776 filed December 30, 1999, and U.S. Application No. 09/492,939 filed January 27, 2000 (each of which applications are hereby incorporated by reference in their entirety).

TECHNICAL FIELD

10 This invention is generally directed to estrogen antagonists and agonists, and to compounds for inhibiting cytokines, and more specifically to compounds which selectively modulate estrogen receptor-beta (ER- β) activity, as well as to pharmaceutical compositions and methods related thereto.

BACKGROUND OF THE INVENTION

15 The estrogen hormone has a broad spectrum of effects on tissues in both females and males. Many of these biological effects are positive, including maintenance of bone density, cardiovascular protection, central nervous system (CNS) function, and the protection of organ systems from the effects of aging. However, in addition to its positive effects, estrogen also is a potent growth factor in
20 the breast and endometrium that increases the risk of cancer.

 Until recently, it was assumed that estrogen binds to a single estrogen receptor (ER) in cells. As discussed below, this simple view changed significantly when a second ER (ER- β) was cloned (with the original ER being renamed ER- α), and when co-factors that modulate the ER response were discovered. Ligands can
25 bind to two different ERs which, in the presence of tissue-specific co-activators and/or co-repressors, bind to an estrogen response element in the regulatory region of genes or to other transcription factors. Given the complexity of ER signaling, along

with the tissue-specific expression of ER- α and ER- β and its co-factors, it is now recognized that ER ligands can act as estrogen agonists and antagonists that mimic the positive effects, or block the negative effects, of estrogen in a tissue-specific manner. This has given rise to the discovery of an entirely new class of drugs, referred to as Selective Estrogen Receptor Modulators or SERMs. These drugs have significant potential for the prevention and/or treatment of cancer and osteoporosis, as well as cardiovascular diseases and neurodegenerative diseases such as Alzheimer's disease.

Bone-resorbing diseases, such as osteoporosis, are debilitating conditions which affect a wide population, and to which there is only limited treatment. For example, osteoporosis affects about 50% of women, and about 10% of men, over the age of 50 in the United States. In individuals with osteoporosis, increased loss of bone mass results in fragile bones and, as a result, increased risk of bone fractures. Other bone-resorption diseases, such as Paget's disease and metastatic bone cancer, present similar symptoms.

Bone is a living tissue which contains several different types of cells. In healthy individuals, the amount of bone made by the osteoblastic cells is balanced by the amount of bone removed or resorbed by the osteoclastic cells. In individuals suffering from a bone-resorbing disease, there is an imbalance in the function of these two types of cells. Perhaps the most well known example of such an imbalance is the rapid increase in bone resorption experienced by postmenopausal women. Such accelerated bone loss is attributed to estrogen deficiency associated with menopause. However, the mechanism of how the loss of estrogen results in increased bone resorption has long been debated.

Recently, investigators have suggested that an increase in bone-resorbing cytokines, such as interleukin-1 (IL-1) and tumor necrosis factor (TNF), may be responsible for postmenopausal bone loss (Kimble et al., *J. Biol. Chem.* 271:28890-28897, 1996), and that inhibitors of these cytokines can partially diminish bone loss following ovariectomy in rodents (Pacifi, *J. Bone Miner Res.* 11:1043-

1051, 1996). Further, discontinuation of estrogen has been reported to lead to an increase in IL-6 secretion by murine bone marrow and bone cells (Girasole et al., *J. Clin. Invest.* 89:883-891, 1992; Jilka et al., *Science* 257:88-91, 1992; Kimble et al., *Endocrinology* 136:3054-3061, 1995; Passeri et al., *Endocrinology* 133:822-828, 5 1993), antibodies against IL-6 can inhibit the increase in osteoclast precursors occurring in estrogen-depleted mice (Girasole et al, *supra*), and bone loss following ovariectomy does not occur in transgenic mice lacking IL-6 (Poli et al., *EMBO J.* 13:1189-1196, 1994).

Existing treatments for slowing bone loss generally involves
10 administration of compounds such as estrogen, bisphosphonates, calcitonin, and raloxifene. These compounds, however, are generally used for long-term treatments, and have undesirable side effects. Further, such treatments are typically directed to the activity of mature osteoclasts, rather than reducing their formation. For example, estrogen induces the apoptosis of osteoclasts, while calcitonin causes the osteoclasts
15 to shrink and detach from the surface of the bone (Hughes et al., *Nat. Med.* 2:1132-1136, 1996; (Jilka et al., *Exp. Hematol.* 23:500-506, 1995). Similarly, bisphosphonates decrease osteoclast activity, change their morphology, and increase the apoptosis of osteoclasts (Parfitt et al., *J. Bone Miner* 11:150-159, 1996; Suzuki et al., *Endocrinology* 137:4685-4690, 1996).

20 Cytokines are also believed to play an important role in a variety of cancers. For example, in the context of prostate cancer, researchers have shown IL-6 to be an autocrine/paracrine growth factor (Seigall et al., *Cancer Res.* 50:7786, 1999), to enhance survival of tumors (Okamoto et al., *Cancer Res.* 57:141-146, 1997), and that neutralizing IL-6 antibodies reduce cell proliferation (Okamoto et al.,
25 *Endocrinology* 138:5071-5073, 1997; Borsellino et al., *Proc. Annu. Meet. Am. Assoc. Cancer Res.* 37:A2801, 1996). Similar results have been reported for IL-6 with regard to multiple myeloma (Martinez-Maza et al., *Res. Immunol.* 143:764-769, 1992; Kawano et al., *Blood* 73:517-526, 1989; Zhang et al., *Blood* 74:11-13, 1989; Garrett et al., *Bone* 20:515-520, 1997; and Klein et al., *Blood* 78:1198-12-4, 1991), renal cell

carcinoma (Koo et al., *Cancer Immunol.* 35:97-105, 1992; Tsukamoto et al., *J. Urol.* 148:1778-1782, 1992; and Weissglas et al., *Endocrinology* 138:1879-1885, 1997), and cervical carcinoma (Estuce et al., *Gynecol. Oncol.* 50:15-19, 1993; Tartour et al., *Cancer Res.* 54:6243-6248, 1994; and Iglesias et al., *Am. J. Pathology* 146:944-952, 5 1995).

Furthermore, IL-6 is also believed to be involved in arthritis, particularly in adjuvant-, collagen- and antigen-induced arthritis (Alonzi et al., *J. Exp. Med.* 187:146-148, 1998; Ohshima et al., *Proc. Natl. Acad. Sci. USA* 95:8222-8226, 1998; and Leisten et al., *Clin. Immunol. Immunopathol* 56:108-115, 1990), and anti- 10 IL-6 antibodies have been reported for treatment of arthritis (Wendling et al., *J. Rheumatol.* 20:259-262, 1993). In addition, estrogen has been shown to induce suppression of experimental autoimmune encephalomyelitis and collagen-induced arthritis in mice (Jansson et al., *Neuroimmunol.* 53:203-207, 1994).

As noted above, it had previously been assumed that estrogen binds to 15 a single estrogen receptor (ER) in cells, causing conformational changes that result in release from heat shock proteins and binding of the receptor as a dimer to the so-called estrogen response element in the promoter region of a variety of genes. Further, pharmacologists have generally believed that non-steroidal small molecule ligands compete for binding of estrogen to ER, acting as either antagonists or agonists 20 in each tissue where the estrogen receptor is expressed. Thus, such ligands have traditionally been classified as either pure antagonists or agonists. This is no longer believed to be correct.

Rather, it is now known that estrogen modulates cellular pharmacology through gene expression, and that the estrogen effect is mediated by 25 estrogen receptors. As noted above, there are currently two estrogen receptors, ER- α and ER- β . The effect of estrogen receptor on gene regulation can be mediated by a direct binding of ER to the estrogen response element (ERE) - "classical pathway" (Jeltsch et al., *Nucleic Acids Res.* 15:1401-1414, 1987; Bodine et al., *Endocrinology* 139:2048-2057, 1998), binding of ER to other transcription factors such as NF- κ B,

C/EBP- β or AP-1 - "non-classical pathway" (Stein et al., *Mol. Cell Biol.* 15:4971-4979, 1995; Paech et al., *Science* 277:1508-1510, 1997; Duan et al., *Endocrinology* 139:1981-1990, 1998), and through non-genomic effects involving ion channel receptors (Watters et al., *Endocrinology* 138:4030-4033, 1997; Improta-Brears et al.,
5 *Proc. Natl. Acad. Sci. USA* 96:4686-4691, 1999; Gu et al., *Endocrinology* 140:660-666, 1999; Beyer et al., *Eur. J. Neurosci.* 10:255-262, 1998).

Progress over the last few years has shown that ER associates with co-activators (e.g., SRC-1, CBP and SRA) and co-repressors (e.g., SMRT and N-CoR), which also modulate the transcriptional activity of ER in a tissue-specific and ligand-
10 specific manner. In addition, evidence now suggests that the majority of estrogen-regulated genes do not have a classical estrogen response element. In such cases, ER interacts with the transcription factors critical for regulation of these genes. Transcription factors known to be modulated in their activity by ER include, for example, AP-1, NF- κ B, C/EBP and Sp-1.

15 Given the complexity of ER signaling, as well as the various types of tissue that express ER and its co-factors, it is now believed that ER ligands can no longer simply be classified as either pure antagonists or agonists. Therefore, the term "selective estrogen receptor modulator" (SERM) has been coined. SERMs bind to ER, but may act as an agonist or antagonist of estrogen in different tissues and on
20 different genes. For example, two of the most well known drugs that behave as SERMs are Tamoxifen and Raloxifene. Studies with these two compounds, as well as other SERMs now in development, have demonstrated that the affinity of a SERM for its receptor in many cases does not correlate with its biological activity. Therefore, ligand-binding assays traditionally used in screening for novel ER
25 modulators have not distinguished between tissue-selectivity and agonist/antagonist behavior.

More recently, a second estrogen receptor, ER- β , has been identified and cloned (Katzenellenbogen and Korach *Endocrinology* 138, 861-2 (1997); Kuiper et al., *Proc. Natl. Acad. Sci. USA* 93, 5925-5930, 1996; Mosselman et al., *FEBS Lett.*

392, 49-53, 1996). ER- β , and the classical ER renamed ER- α , have significantly different amino acid sequences in the ligand binding domain and carboxy-terminal transactivation domains (~56% amino acid identity), and only 20% homology in their amino-terminal transactivation domain. This suggests that some ligands may have higher affinity to one receptor over the other. Further, ligand-dependent conformational changes of the two receptors, and interaction with co-factors, will result in very different biological actions of a single ligand. In other words, a ligand that acts as an agonist on ER- α may very well serve as an antagonist on ER- β . An example of such behavior has been described by Paech et al. (*Science* 277, 1508-1510, 1997). In that paper, estrogen is reported to activate an AP-1 site in the presence of ER- α , but to inhibit the same site in the presence of ER- β . In contrast, Raloxifene (Eli Lilly & Co.) and Tamoxifen and ICI-182,780 (Zeneca Pharmaceuticals) stimulate the AP-1 site through ER- β , but inhibit this site in the presence of ER- α . Another example has been described by Sun et al. (*Endocrinology* 140, 800-4, 1999). In this paper, the R,R-enantiomer of a tetrahydrochrysene is reported to be an agonist on ER- α , but a complete antagonist on ER- β , while the S,S-enantiomer is an agonist on both receptors.

Furthermore, ER- α and ER- β have both overlapping and different tissue distributions, as analyzed predominantly by RT-PCR or *in-situ* hybridization due to a lack of good ER- β antibodies. Some of these results, however, are controversial, which may be attributable to the method used for measuring ER, the species analyzed (rat, mouse, human) and/or the differentiation state of isolated primary cells. Very often tissues express both ER- α and ER- β , but the receptors are localized in different cell types. In addition, some tissues (such as kidney) contain exclusively ER- α , while other tissues (such as uterus, pituitary and epidymis) show a great predominance of ER- α (Couse et al., *Endocrinology* 138, 4613-4621, 1997; Kuiper et al., *Endocrinology* 138, 863-870, 1997). In contrast, tissues expressing high levels of ER- β include prostate, testis, ovaries and certain areas of the brain (Brandenberger et al., *J. Clin. Endocrinol. Metab.* 83, 1025-8, 1998; Enmark et al., *J.*

Clinic. Endocrinol. Metabol. 82, 4258-4265, 1997; Laflamme et al., *J. Neurobiol.* 36, 357-78, 1998; Sar and Welsch, *Endocrinology* 140, 963-71, 1999; Shughrue et al., *Endocrinology* 138, 5649-52, 1997a; Shughrue et al., *J. Comp. Neurol.* 388, 507-25, 1997b).

- 5 The development of ER- α (Korach, *Science* 266, 1524-1527, 1994) and ER- β (Krege et al., *Proc. Natl. Acad. Sci. USA* 95, 15677-82, 1998) knockout mice further demonstrate that ER- β has different functions in different tissues. For example, ER- α knockout mice (male and female) are infertile, females do not display sexual receptivity and males do not have typical male-aggressive behavior (Cooke et
- 10 al., *Biol. Reprod.* 59, 470-5, 1998; Das et al., *Proc. Natl. Acad. Sci. USA* 94, 12786-12791, 1997; Korach, 1994; Ogawa et al., *Proc. Natl. Acad. Sci. USA* 94, 1476-81, 1997; Rissman et al., *Endocrinology* 138, 507-10, 1997a; Rissman et al., *Horm. Behav.* 31, 232-243, 1997b). Further, the brains of these animals still respond to estrogen in a pattern that is similar to that of wild type animals (Shughrue et al., *Proc.*
- 15 *Natl. Acad. Sci. USA* 94, 11008-12, 1997c), and estrogen still inhibits vascular injury caused by mechanical damage (Iafrati et al., *Nature Med.* 3, 545-8, 1997). In contrast, mice lacking the ER- β develop normally, are fertile and exhibit normal sexual behavior, but have fewer and smaller litters than wild-type mice (Krege et al., 1998), have normal breast development and lactate normally. The reduction in
- 20 fertility is believed to be the result of reduced ovarian efficiency, and ER- β is the predominant form of ER in the ovary, being localized in the granulosa cells of maturing follicles.


- In summary, compounds which serve as estrogen antagonists or agonists have long been recognized for their significant pharmaceutical utility in the
- 25 treatment of a wide variety of estrogen-related conditions, including conditions related to the brain, bone, cardiovascular system, skin, hair follicles, immune system, bladder and prostate (Barkhem et al., *Mol. Pharmacol.* 54, 105-12, 1998; Farhat et al., *FASEB J.* 10, 615-624, 1996; Gustafsson, *Chem. Biol.* 2, 508-11, 1998; Sun et al., 1999; Tremblay et al., *Endocrinology* 139, 111-118, 1998; Turner et al.,

Endocrinology 139, 3712-20, 1998). In addition, a variety of breast and non-breast cancer cells have been described to express ER, and serve as the target tissue for specific estrogen antagonists (Brandenberger et al., 1998; Clinton and Hua, *Crit. Rev. Oncol. Hematol.* 25, 1-9, 1997; Hata et al., *Oncology* 55 Suppl 1, 35-44, 1998; Rohlf et al., *Prostate* 37, 51-9, 1998; Simpson et al., *J Steroid Biochem Mol Biol* 64, 137-45, 1998; Yamashita et al., *Oncology* 55 Suppl 1, 17-22, 1998).

With the recent identification of the ER- β , and the recognition that ER- α and ER- β have different biological roles, ER-selective modulators would similarly possess significant clinical utility. Since ER- β is expressed strongly in a number of tissues including prostate, bladder, ovary, testis, lung, small intestine, vascular endothelium, and various parts of the brain, compounds that selectively modulate ER- β would be of clinical importance in the treatment of a variety of diseases or conditions, such as prostate cancer, testicular cancer, cardiovascular diseases, neurodegenerative disorders, urinary incontinence, CNS, GI tract conditions, and bone and other cancers. Such compounds would have minimal effect on tissues that contains ER- α , and thus exhibit different side-effect profiles. For example, while estrogen replacement therapy is associated with a variety of beneficial effects (such as bone protection, cardiovascular effect, prevention of hot flashes, dementia, bone metabolism, etc.), such therapy also has adverse effects (such as breast and endometrial cancer, thrombosis, etc.). Some of these adverse effects are believed to be mediated by ER- α or ER- β specific mechanisms. Thus, ER- β antagonists or agonists will display different therapeutic profiles compared to ER- α antagonists or agonists, and would be preferentially beneficial in tissues expressing high levels of ER- β (see, e.g., Nilsson et al., *TEM* 9, 387-395, 1998; Chang and Prins, *The Prostate* 40, 115-124, 1999). Furthermore, a number of investigators have shown that environmental chemicals and phytoestrogens preferentially interact with ER- β by triggering biological responses similar to that of estrogen (see, e.g., Kuiper et al., *Endocrinology* 139, 4252-4263, 1998). Thus, compounds that antagonize ER- β

would also be important in regulating interactions with chemicals, affecting health, reproductive capacity and the like.

In recent years a number of both steroidal and nonsteroidal compounds which interact with ER have been developed. For example, Tamoxifen was originally
5 developed as an anti-estrogen and used for the treatment of breast cancer, but more recently has been found to act as a partial estrogen agonist in the uterus, bone and cardiovascular system. Raloxifene is another compound that has been proposed as a SERM, and has been approved for treatment of osteoporosis.



Analogues of Raloxifene have also been reported (Grese et al., *J. Med. Chem.* 40:146-167, 1997).

15 As for coumarin-based compounds, a number of structures have been proposed, including the following: Roa et al., *Synthesis* 887-888, 1981; Buu-Hoi et al., *J. Org. Chem.* 19:1548-1552, 1954; Gupta et al., *Indian J. Exp. Biol.* 23:638-640, 1985; Published PCT Application No. WO 96/31206; Verma et al., *Indian J. Chem.* 32B:239-243, 1993; Lednicer et al., *J. Med. Chem.* 8:725-726, 1965; Micheli et al.,
20 *Steroids* 5:321-335, 1962; Brandt et al., *Int. J. Quantum Chemistry: Quantum Biol. Symposia* 13:155-165, 1986; Wani et al., *J. Med. Chem.* 18:982-985, 1975; Pollard et al., *Steroids* 11:897-907, 1968.


Accordingly, there is a need in the art for estrogen antagonists and agonists generally, and more specifically for compounds that inhibit cytokines, particularly IL-6, including pharmaceutical compositions containing such compounds as well as methods relating to the use thereof. There is also a need for compounds
5 that selectively modulates ER- β . The present invention fulfills these needs, and provides other related advantages.

SUMMARY OF THE INVENTION

In brief, the present invention is generally directed to estrogen antagonists and/or agonists, including pharmaceutical compositions containing the
10 same, as well as to methods for treating estrogen-related conditions. Such conditions are more specifically discussed below, and generally include (but are not limited to) obesity, breast cancer, osteoporosis, endometriosis, cardiovascular disease, prostate cancer, menopausal syndromes, hair loss (alopecia), type-II diabetes, Alzheimer's disease, urinary incontinence, GI tract conditions, spermatogenesis, vascular
15 protection after injury, endometriosis, learning and memory, CNS effects, plasma lipid levels, acne, cataracts, hirsutism, other solid cancers (such as colon, lung, ovarian, melanoma, CNS, and renal), multiple myeloma, and lymphoma.

In a more specific embodiment, this invention is directed to compounds for inhibiting cytokines such as interleukin-6 (IL-6) and to methods
20 relating to the treatment of conditions associated therewith, as well as pharmaceutical compositions containing one or more of the compounds of this invention. In this context, treating conditions associated with increased levels of cytokines include (but are not limited to) methods for treating cancer, arthritis and bone-resorbing diseases, particularly for reducing formation of osteoclasts and/or blocking cytokine production
25 in the context of osteoporosis.

The compounds of this invention have the following general structure (I):



(I)

wherein R₁, R₂, R₃, *n* and *p* are as defined in the following detailed description,
 5 including stereoisomers, prodrugs and pharmaceutically acceptable salts thereof.

As noted above, the compounds of this invention have utility over a wide range of therapeutic and prophylactic applications, and may be used to treat a variety of diseases associated with bone resorption, as well as for treatment of cancer and arthritis. Such methods involve the administration of an effective amount of a
 10 compound of this invention, preferably in the form of a pharmaceutical composition, to an animal in need thereof (including a human).

In another embodiment, methods are disclosed for modulating cells and/or tissues that express ER-β by contacting the cell and/or tissue with an effective amount of a compound of structure (I). In an embodiment, the cell and/or tissue
 15 preferentially expresses ER-β over ER-α, such as cell and/or tissue of bone, bladder, uterus, ovary, prostate, testis, epididymis, gastrointestinal (GI) tract, kidney, breast, heart, vessel wall, immune system, lung, pituitary, hippocampus and hypothalamus.

In still a further embodiment, the present invention discloses methods for treating an estrogen-related condition by administering to an warm-blooded
 20 animal in need thereof an effective amount of a compound of structure (I) formulated as a pharmaceutical composition suitable for administration to the animal. In representative embodiments, the estrogen-related condition is breast cancer, osteoporosis, endometriosis, cardiovascular disease, hypercholesterolemia, prostatic

hypertrophy, obesity, prostate cancer, menopausal syndromes, type-II diabetes, Alzheimer's disease, urinary incontinence, GI tract conditions, spermatogenesis, vascular protection after injury, endometriosis, learning and memory, CNS effects, plasma lipid levels, acne, hirsutism, other solid cancers (such as colon, lung, ovarian, testis, melanoma, CNS, and renal), multiple myeloma, lymphoma, prostatic carcinomas, obesity, hot flashes, skin effects, mood swings, memory loss, and/or adverse reproductive effects associated with exposure to environmental chemicals. In other embodiments, methods are disclosed for inhibiting a cytokine, such as IL-6 and GM-CSF, in an animal in need thereof, as well as methods for treating cancer associated therewith.

These and other aspects of this invention will be evident upon reference to the following detailed description and attached drawings. To that end, various references are set forth herein which describe in more detail certain aspects of this invention, and which are hereby incorporated by reference in their entirety.

BRIEF DESCRIPTION OF THE DRAWINGS

Figures 1A and 1B illustrate the activity of a representative compound of this invention to inhibit production of IL-6 and GM-CSF, respectively.

Figure 2 illustrates the activity of a representative compound of this invention to inhibit IL-6 production in rheumatoid arthritis synoviocytes.

Figure 3 illustrates the activity of a representative compound of this invention to inhibit breast cancer proliferation.

Figure 4 illustrates the activity of a representative compound of this invention to inhibit a prostate cancer cell line.

DETAILED DESCRIPTION OF THE INVENTION

As mentioned above, the present invention is directed to compounds that have activity as estrogen antagonists and/or agonists, as well as pharmaceutical compositions containing one or more of such compounds and methods related to the


same. As estrogen antagonists and/or agonists, the compounds of this invention have utility in the treatment of a wide range of estrogen-related conditions. In this context, the term "treatment" includes both treatment and/or prevention of an estrogen-related condition. Thus, the compounds of this invention may be administered as a
5 therapeutic and/or prophylactic agent. An estrogen "agonist" is a compound that binds to ER and mimics the action of estrogen in one or more tissues, while an "antagonist" binds to ER and blocks the action of estrogen in one or more tissues. Further, the term "estrogen-related condition" encompasses any condition associated with elevated or depressed levels of estrogen, a selective estrogen receptor
10 modulators (SERM), ER- α or ER- β .

Accordingly, the compounds of the present invention may also be used within a method for treating estrogen-related conditions, including (but not limited to) breast cancer, osteoporosis, endometriosis, cardiovascular disease, hypercholesterolemia, prostatic hypertrophy, prostatic carcinomas, obesity, hot
15 flashes, skin effects, mood swings, memory loss, menopausal syndromes, hair loss (alopecia), type-II diabetes, Alzheimer's disease, urinary incontinence, GI tract conditions, spermatogenesis, vascular protection after injury, endometriosis, learning and memory, CNS effects, plasma lipid levels, acne, hirsutism, other solid cancers (such as colon, lung, ovarian, testis, melanoma, CNS, and renal), multiple myeloma,
20 cataracts, lymphoma, and adverse reproductive effects associated with exposure to environmental chemicals.

In one aspect of this invention, it has been found that the compounds of this invention are selective estrogen receptor modulators and, more specifically, are selective to ER- β . Accordingly, in this aspect of the invention, the compounds
25 have utility as agents for treatment of osteoporosis, hormonally regulated cancer, combination treatment of cancer (such as with taxol cisplatin or chemotherapy), women's gynecologic health issues, and health conditions resulting from exposure to environmental chemicals.

The compounds of this invention have the following general structure

(I):



including stereoisomers, prodrugs and pharmaceutically acceptable salts thereof,
wherein:


n is 0, 1, 2, 3 or 4;

p is 0, 1 or 2;

R_1 is an unsubstituted or substituted C_{6-12} aryl, C_{7-12} arylalkyl, C_{3-12} heterocycle or C_{4-16} heterocyclealkyl;

R_2 is NR_aR_b wherein R_a and R_b are independently hydrogen, C_{1-8} alkyl, C_{6-12} aryl, or heterocycle, and wherein R_a and R_b are optionally substituted with up to three substituents independently selected from C_{1-6} alkyl, halogen, C_{1-6} alkoxy, hydroxy and carboxyl;

or R_2 is a heterocyclic ring of the following structure:



wherein

m_1 and m_2 are independently 0, 1 or 2, and both of m_1 and m_2 are not 0,

A is CH₂, O, S or NH,

Z represents 0, 1, 2 or 3 heterocyclic ring substituents selected from halogen, C₁₋₈alkyl, C₆₋₁₂aryl, C₇₋₁₂arylalkyl, C₃₋₁₂heterocycle, or C₄₋₁₆heterocyclealkyl,

and wherein any hydrogen atom on the heterocyclic ring may, taken together with a hydrogen atom on an adjacent atom of the heterocyclic ring, form a double bond;

R₃ is hydrogen, R₄, C(=O)R₄, C(=O)OR₄, CONHR₄, CONR₄R₅, or SO₂NR₅R₅;

R₄ and R₅ are independently C₁₋₈alkyl, C₆₋₁₂aryl, C₇₋₁₂arylalkyl, or a five- or six-membered heterocycle containing up to two heteroatoms selected from O, NR₆ and S(O)_q, wherein each of the above groups are optionally substituted with one to three substituents independently selected from R₇ and q is 0, 1 or 2;

R₆ is hydrogen or C₁₋₄alkyl; and

R₇ is hydrogen, halogen, hydroxy, C₁₋₆alkyl, C₁₋₄alkoxy, C₁₋₄acyloxy, C₁₋₄thio, C₁₋₄alkylsulfinyl, C₁₋₄alkylsulfonyl, (hydroxy)C₁₋₄alkyl, C₆₋₁₂aryl, C₇₋₁₂arylalkyl, COOH, CN, CONHOR₈, SO₂NHR₈, NH₂, C₁₋₄alkylamino, C₁₋₄dialkylamino, NHSO₂R₈, NO₂, or a five- or six-membered heterocycle, where each occurrence of R₈ is independently C₁₋₆alkyl.

As used herein, a "C₆₋₁₂aryl" is an aromatic moiety containing from 6 to 12 carbon atoms. In one embodiment, the C₆₋₁₂aryl is selected from (but not limited to) phenyl, tetralinyl, and naphthalenyl.

A "C₇₋₁₂aralkyl" is an arene containing from 7 to 12 carbon atoms, and has both aliphatic and aromatic units. In one embodiment, the C₇₋₁₂aralkyl is selected from (but not limited to) benzyl, ethylbenzyl (*i.e.*, -(CH₂)₂phenyl), propylbenzyl and isobutylbenzyl.

- 5 A "C₃₋₁₂heterocycle" is a compound that contains a ring made up of more than one kind of atom, and which contains 3 to 12 carbon atoms, including (but not limited to) pyrrolyl, furanyl, thienyl, imidazolyl, oxazolyl, thiazolyl, pyrazolyl, pyrrolidinyl, pyridinyl, pyrimidinyl and purinyl.

- A "C₄₋₁₆heterocyclealkyl" is a compound that contains a
10 C₃₋₁₂heterocycle linked to a C₁₋₈alkyl.



A "C₁₋₈alkyl" is a straight chain or branched carbon chain containing from 1 to 8 carbon atoms, including (but not limited to) methyl, ethyl, and n-propyl. Similarly, a "C_{1-x}alkyl" has the same meaning, but wherein "x" represents the number of carbon atoms less than eight, such as C₁₋₆alkyl.

- 15 A "substituted" C_{1-x}alkyl, C₆₋₁₂aryl, C₇₋₁₂aralkyl, C₃₋₁₂heterocycle, or C₄₋₁₆heterocyclealkyl moiety is a C_{1-x}alkyl, C₆₋₁₂aryl, C₇₋₁₂aralkyl, C₃₋₁₂heterocycle, or C₄₋₁₆heterocyclealkyl moiety having at least one hydrogen atom replaced with a substituent.

- A "substituent" is a moiety selected from halogen, -OH, -R', -OR',
20 -COOH, -COOR', -COR', -CONH₂, -NH₂, -NHR', -NR'R', -SH, -SR', -SOOR', -SOOH and -SOR', where each occurrence of R' is independently selected from an unsubstituted or substituted C₁₋₈alkyl, C₆₋₁₂aryl, C₇₋₁₂aralkyl, C₃₋₁₂heterocycle or C₄₋₁₆heterocyclealkyl.





A "halogen" is fluoro, chloro, bromo or iodo.

- 25 In one embodiment of this invention, A of the heterocyclic ring R₂ is CH₂, m₁ is 1, and m₂ is 0 or 1, as represented by the following structures (i) and (ii), respectively:





In structures (i) and (ii) above, it should be noted that the hydrogen atoms are not depicted in order to clarify that the optional Z substituent(s) may be on any atom of the heterocyclic ring, and that the point of attachment to structure (I) may be through a carbon or nitrogen atom.

Thus, in more specific embodiments of structures (i) and (ii) wherein Z is present, R_2 includes the following structures (iii) through (vi):



wherein Z is, for example, methyl.


In another embodiment, A of the heterocyclic ring R_2 is O or NH, m_1 is 1, and m_2 is 0 or 1, as represented by, for example, the following structures (vii) and (viii):



As with structures (i) and (ii) above, in structures (vii) and (viii) the hydrogen atoms are not depicted in order to clarify that the optional Z substituent(s) may be on any


atom of the heterocyclic ring, and that the point of attachment to structure (I) may be through a carbon or nitrogen atom.

In addition to the above depicted structures, any hydrogen atom of the heterocyclic ring may be taken together with a hydrogen atom attached to an adjacent
 5 heterocyclic ring atom to form a double bond. For example, with regard to structure (vii) above, corresponding unsaturated analogs include the following structures (ix), (x) and (xi):



In one embodiment of this invention, R_1 is an unsubstituted or substituted phenyl, and the compounds of this invention have the following structure (II):


15



wherein X represents one or more optional substituents as defined above, and R_2 , R_3 , n and p are as defined above.


20

In another embodiment, R_3 is hydrogen, as represented by structure (III):



wherein R_1 , R_2 , n and p are as defined above.

- 5 In a more specific embodiments of structures (II) and (III), representative compounds of this invention have the following structure (IV):




10

wherein A , X , Z , m_1 , m_2 , n and p are as defined above.

In a further embodiment of structure (IV), m_1 , m_2 and p are 1, A is CH_2 , the optional Z substituent is not present, and n is 2, and compounds of this invention have the following structure (V):


15

20



wherein X represents one or more optional substituents as defined above.


In a more specific embodiment, X is either (a) not present or (b)
 5 represents a single substituent, such as a single substituent at the para position.
 Accordingly, representative compounds of this invention include (but are not limited
 to) compounds having the following structures (VIa) and (VIb):



wherein X in structure (VIb) represents a halogen, such as fluorine or chlorine.

In a further embodiment of this invention, p is 0, and compounds of
 this invention have the following structure (VII):


15



wherein R_1 , R_2 , R_3 and n are as defined above.

In one embodiment of structure (VII), R_1 is an unsubstituted or substituted phenyl, and the compounds of this invention have the following structure (VIII):

5




(VIII)

wherein X represents one or more optional substituents as defined above, and R_2 , R_3 and n are as defined above.

10


In another embodiment, R_3 is hydrogen, as represented by structure (IX):



(IX)


15 wherein R_1 , R_2 and n are as defined above.

In a more specific embodiments of structures (VIII) and (IX), representative compounds of this invention have the following structure (X):





wherein A, X, Z, m_1 , m_2 , and n are as defined above.

- 5 In a further embodiment of structure (X), m_1 and m_2 are 1, A is CH_2 , the optional Z substituent is not present, and n is 2, and compounds of this invention have the following structure (XI):



10 wherein X represents one or more optional substituents as defined above.

- In a more specific embodiment, X is either (a) not present or (b) represents a single substituent, such as a single substituent at the para position. Accordingly, representative compounds of this invention include (but are not limited to) compounds having the following structures (XIa) and (XIb):
- 15





- 5 wherein X in structure (XIb) represents a halogen, such as fluorine or chlorine.

In the embodiment wherein p is 0, particularly when R_1 is hydrogen and the phenyl group at the 4-position of the coumarin ring is para-substituted, the compounds of this invention have been found to exhibit selectivity for ER- β over ER- α .

- 10 The compounds of this invention may be made by one skilled in organic synthesis by known techniques, as well as by the synthetic routes disclosed herein. For example, representative compounds of this invention may be synthesized by the following general Reaction Scheme 1:


Reaction Scheme 1





Reaction Scheme 1 yields compounds wherein R_3 is methyl or
 5 hydrogen, and R_2 is a heterocyclic ring as defined in structure (I). Further
 substitution at the R_3 position may be accomplished using an appropriately
 substituted phenol, or by subsequent conversion of the hydroxyl group (when $R_3 = H$)
 using techniques known in the field of organic synthesis. Similarly, compounds of
 structure (I) wherein R_2 is NR_4R_5 may be made by employing the corresponding
 10 amino chloride, $R_4R_5N(CH_2)_nCl$, in place of the heterocyclic ring in the second-to-last
 step of Reaction Scheme 1.

More specifically, representative compounds of this invention (when
 R_3 is hydrogen and R_2 is piperid-1-yl) may be made by the following Reaction
 Scheme 2:

Reaction Scheme 2

5

With regard to stereoisomers, the compounds of structure (I) may have chiral centers and may occur as racemates, racemic mixtures and as individual enantiomers or diastereomers. All such isomeric forms are included within the present invention, including mixtures thereof. Furthermore, some of the crystalline forms of the compounds of structure (I) may exist as polymorphs, which are included in the present invention. In addition, some of the compounds of structure (I) may also form solvates with water or other organic solvents. Such solvates are similarly included within the scope of this invention.

Although not intending to be limited by the following theory, in the context of bone-resorbing diseases it is believed that the compounds of this invention function by blocking cytokine production and/or by inhibiting formation of osteoclasts. The cytokine IL-6 has previously been shown to be an important factor in inducing the formation of osteoclasts (Girasole et al., *supra*; Jilka et al. (1992), *supra*; Jilka et al. (1995), *supra*; Kimble et al. (1995), *supra*; Pacifici et al., *supra*; and Passeri et al., *supra*). Other investigators have shown that administration of the neutralizing antibody, antisense oligos, or the Sant 5 antagonist against IL-6, reduces the number of osteoclasts in trabecular bone of ovariectomized mice (Devlin et al., *J. Bone Miner* 13:393-399, 1998; Girasole et al., *supra*; Jilka et al. (1992), *supra*; and Schiller et al., *Endocrinology* 138:4567-4571, 1997), the ability of human giant cells to resorb dentine (Ohsaki et al., *Endocrinology* 131:2229-2234, 1993; and Reddy et al., *J. Bone Min. Res.* 9:753-757, 1994), and the formation of osteoclasts in normal human bone marrow culture. It has also been found that estrogen downregulates the IL-6 promoter activity by interactions between the estrogen receptor and the transcription factors NF- κ B and C/EBP β (Stein et al., *Mol. Cell Biol.* 15:4971-4979, 1995).

Granulocyte-macrophage colony-stimulating factor (GM-CSF) has been suggested to play a role in the proliferation of osteoclastic precursor cells. In long term cultures of human or mouse bone marrow cells or peripheral blood cells, GM-CSF promotes the formation of osteoclastic cells (Kurihara et al., *Blood* 74:1295-1302, 1989; Lorenzo et al., *J. Clin. Invest.* 80:160-164, 1987; MacDonald et al., *J. Bone Miner* 1:227-233, 1986; and Shinar et al., *Endocrinology* 126:1728-1735, 1990). Bone marrow cells isolated from postmenopausal women, or women who discontinued estrogen therapy, expressed higher levels of GM-CSF than cells from premenopausal women (Bismar et al., *J. Clin. Endocrinol. Metab.* 80:3351-3355, 1995). Expression of GM-CSF has also been shown to be associated with the tissue distribution of bone-resorbing osteoclasts in patients with erosion of orthopedic implants (Al-Saffar et al., *Anatomic Pathology* 105:628-693, 1996).

In addition, it has been found that blocking the action of IL-6 with an anti-IL-6 antibody reduces the formation of osteoclast-like cells in the human bone cell co-culture system of Example 8. This system includes immortalized human osteoblasts and premonocytic cell line U937, which are co-cultured in the same tissue culture well. Under appropriate culture conditions the osteoblasts secrete IL-6 which acts on the premonocytic cells causing their differentiation into osteoclast-like cells. Thus, the co-culture system more closely reflects the physiological environment of bone, and provides a functional readout for the efficacy of the compounds known to block the production of IL-6 and leading to osteoclast activation and formation. For example, estrogen was found to suppress the production of IL-6 in the co-culture system of Example 8.

Furthermore, an anti-GM-CSF antibody was found to reduce the formation of osteoclast-like cells, and estrogen reduced the production of GM-CSF in the co-culture system of Example 8. This demonstrates that estrogen may exert its inhibitory effect on osteoclast formation and function through blocking the production of IL-6 and GM-CSF. Therefore, compounds of this invention that block IL-6 and/or GM-CSF production, or inhibit formation and function of osteoclasts, or both, are useful in the treatment of bone-resorbing diseases.

As mentioned above, cytokines are also believed to play an important role in a variety of cancers. In the context of prostate cancer, researchers have shown IL-6 to be an autocrine/paracrine growth factor (Seigall et al., *supra*), to enhance survival of tumors (Okamoto et al. (*Cancer Res.* 1997), *supra*), and that neutralizing IL-6 antibodies reduce cell proliferation (Okamoto et al. (*Endocrinology* 1997), *supra*; Borsellino et al., *supra*). IL-6 has also been reported to play a role with regard to multiple myeloma (Martinez-Maza et al., *supra*; Kawano et al., *supra*; Zhang et al., *supra*; Garrett et al., *supra*; and Klein et al., *supra*), renal cell carcinoma (Koo et al., *supra*; Tsukamoto et al., *supra*; and Weissglas et al., *supra*), and cervical carcinoma (Estuce et al., *supra*; Tartour et al., *supra*; and Iglesias et al., *supra*).

IL-6 is also believed to be involved in arthritis, particularly in adjuvant-, collagen- and antigen-induced arthritis (Alonzi et al., *supra*; Ohshima et al., *supra*; and Leisten et al., *supra*), and anti-IL-6 antibodies have been reported for treatment of arthritis (Wendling et al., *supra*). In addition, estrogen has been shown
5 to induce suppression of experimental autoimmune encephalomyelitis and collagen-induced arthritis in mice (Jansson et al., *supra*).

Accordingly, other embodiments of the present invention include methods for treating bone-resorbing diseases generally, including (but not limited to) osteoporosis, metastatic bone cancer and hypercalcemia, osteolytic lesions with
10 orthopedic implants, Paget's disease, and bone loss associated with hyperparathyroidism. Other conditions associated with IL-6 include various cancers and arthritis. Representative cancers are breast cancer, prostate cancer, colon cancer, endometrial cancer, multiple myeloma, renal cell carcinoma, and cervical carcinoma. Arthritic conditions include adjuvant-, collagen-, bacterial- and antigen-induced
15 arthritis, particularly rheumatoid arthritis.

In addition, the compounds of the present invention also act as estrogen antagonists and/or agonists, and have utility in the treatment of a wide range of estrogen-related conditions. In this context, treatment includes both treatment and/or prevention of an estrogen-related condition. Thus, the compounds of this
20 invention may be administered as a therapeutic and/or prophylactic agent. An estrogen "agonist" is a compound that binds to ER and mimics the action of estrogen in one or more tissues, while an "antagonist" binds to ER and blocks the action of estrogen in one or more tissues. Further, the term "estrogen-related condition" encompasses any condition associated with elevated or depressed levels of estrogen, a
25 selective estrogen receptor modulators (SERM) or ER. In this context, ER includes both ER- α and/or ER- β , as well as any isoforms, mutations and proteins with significant homology to ER.

Accordingly, the compounds of the present invention may also be used within a method for treating estrogen-related conditions, including (but not limited to)

breast cancer, osteoporosis, endometriosis, cardiovascular disease, hypercholesterolemia, prostatic hypertrophy, prostatic carcinomas, obesity, hot flashes, skin effects, mood swings, memory loss, prostate cancer, menopausal syndromes, hair loss (alopecia), type-II diabetes, Alzheimer's disease, urinary
5 incontinence, GI tract conditions, spermatogenesis, vascular protection after injury, endometriosis, learning and memory, CNS effects, plasma lipid levels, acne, cataracts, hirsutism, other solid cancers (such as colon, lung, ovarian, melanoma, CNS, and renal), multiple myeloma, lymphoma, and adverse reproductive effects associated with exposure to environmental chemicals or natural hormonal imbalances.

10 Compounds of the present invention that are estrogen agonists are useful for oral contraception; relief for the symptoms of menopause; prevention of threatened or habitual abortion; relief of dysmenorrhea; relief of dysfunctional uterine bleeding; relief of endometriosis; an aid in ovarian development; treatment of acne; diminution of excessive growth of body hair in women (hirsutism); the prevention
15 and treatment of cardiovascular disease; prevention and treatment of atherosclerosis; prevention and treatment of osteoporosis; treatment of benign prostatic hyperplasia and prostatic carcinoma obesity; and suppression of post-partum lactation. These agents also have a beneficial effect on plasma lipid levels and as such are useful in treating and preventing hypercholesterolemia. Those which are estrogen antagonists
20 are useful as antiestrogens in, for example, breast and ovarian tissue and thus are useful in the treatment and prevention of breast and ovarian cancer.

 Methods of this invention involve administering a compound of structure (I), or a pharmaceutical composition containing the one or more of the same, to an animal in need thereof in an amount sufficient to treat the disease or condition
25 of interest. To that end, the term "treat" (or the related terms "treating" and "treatment") means administration of a compound, typically in combination with an appropriate delivery vehicle or agent, to an animal that does not show signs of a disease or condition (e.g., prophylactic or preventative administration) or that does show signs of a disease or condition (e.g., curative or treatment administration).

Further, the phrase "effective amount" means a dose of compound that, after a given time, results in the desired effect. For example, in the context of bone-resorbing disease, an effective amount results in bones mass that is statistically different from that of animals treated with placebo. Similarly, for cancer and arthritis, an effective
5 amount is an amount sufficient to produce the desired effect on the cancerous or arthritic tissue.

The methods of this invention include administration of an effective amount of a compound of structure (I), or a salt thereof, as the active ingredient. Pharmaceutically acceptable salts of the compounds of structure (I) are typically salts
10 of non-toxic type commonly used, such as salts with organic acids (e.g., formic, acetic, trifluoroacetic, citric, maleic, tartaric, methanesulfonic, benzenesulfonic or toluenesulfonic acids), inorganic acids (e.g., hydrochloric, hydrobromic, sulfuric or phosphoric acids), and amino acids (e.g., aspartic or glutamic acids). These salts may be prepared by the methods known to chemists of ordinary skill.

15 The compounds of this invention may be administered to animals (including humans) orally or parenterally in the conventional form of preparations, such as capsules, microcapsules, tablets, granules, powder, troches, pills, suppositories, injections, suspensions and syrups. Suitable formulations may be prepared by methods commonly employed using conventional, organic or inorganic
20 additives, such as an excipient (e.g., sucrose, starch, mannitol, sorbitol, lactose, glucose, cellulose, talc, calcium phosphate or calcium carbonate), a binder (e.g., cellulose, methylcellulose, hydroxymethylcellulose, polypropylpyrrolidone, polyvinylpyrrolidone, gelatin, gum arabic, polyethyleneglycol, sucrose or starch), a disintegrator (e.g., starch, carboxymethylcellulose, hydroxypropylstarch, low
25 substituted hydroxypropylcellulose, sodium bicarbonate, calcium phosphate or calcium citrate), a lubricant (e.g., magnesium stearate, light anhydrous silicic acid, talc or sodium lauryl sulfate), a flavoring agent (e.g., citric acid, menthol, glycine or orange powder), a preservative (e.g., sodium benzoate, sodium bisulfite, methylparaben or propylparaben), a stabilizer (e.g., citric acid, sodium citrate or

acetic acid), a suspending agent (e.g., methylcellulose, polyvinyl pyrrolidone or aluminum stearate), a dispersing agent (e.g., hydroxypropylmethylcellulose), a diluent (e.g., water), and base wax (e.g., cocoa butter, white petrolatum or polyethylene glycol). The amount of the active ingredient in the medical composition
5 may be at a level that will exercise the desired therapeutical effect; for example, about 0.1 mg to 100 mg in unit dosage for both oral and parenteral administration.

The active ingredient may be usually administered one to four times a day with a unit dosage of 0.1 mg to 100 mg in human patients, but the above dosage may be properly varied depending on the age, body weight and medical condition of
10 the patient and the type of administration. A preferred dose is 0.25 mg to 25 mg in human patients. One dose per day is preferred.

Pharmaceutical chemists will recognize that physiologically active compounds which one or more accessible hydroxy groups are frequently administered in the form of pharmaceutically acceptable esters. The literature concerning such
15 compounds, such as estradiol, provides a number of instances of such esters. It is believed that such esters are metabolically cleaved in the body, and that the actual drug is the hydroxy compound itself. It is known known in pharmaceutical arts to adjust the rate or duration of action of a compound by appropriate choices of ester groups.

To this end, prodrugs are also included within the context of this
20 invention. Prodrugs are any covalently bonded carriers that release a compound of structure (I) *in vivo* when such prodrug is administered to a patient. Prodrugs are generally prepared by modifying functional groups in a way such that the modification is cleaved, either by routine manipulation or *in vivo*, yielding the parent
25 compound. Prodrugs include, for example, compounds of this invention wherein the hydroxy group (when $R_3 = H$) is bonded to any group that, when administered to a patient, cleaves to form the hydroxy group.

The pharmaceutically acceptable acid addition salts of the compounds of this invention may be formed of the compound itself, or of any of its esters, and

include the pharmaceutically acceptable salts which are often used in pharmaceutical chemistry. For example, salts may be formed with inorganic or organic acids such as hydrochloric acid, hydrobromic acid, hydroiodic acid, sulfonic acids including such agents as naphthalenesulfonic, methanesulfonic and toluenesulfonic acids, sulfuric acid, nitric acid, phosphoric acid, tartaric acid, pyrosulfuric acid, metaphosphoric acid, succinic acid, formic acid, phthalic acid, lactic acid and the like, most preferable with hydrochloric acid, citric acid, benzoic acid, maleic acid, acetic acid and propionic acid. It is usually preferred to administer a compound of this invention in the form of an acid addition salt, as it is customary in the administration of pharmaceuticals bearing a basic group.

The compounds of this invention, as discussed above, are very often administered in the form of acid addition salts. The salts are conveniently formed, as is usual in organic chemistry, by reacting the compound of this invention with a suitable acid, such as have been described above. The salts are quickly formed in high yields at moderate temperatures, and often are prepared by merely isolating the compound from a suitable acidic wash in the final step of the synthesis. The salt-forming acid is dissolved in an appropriate organic solvent, or aqueous organic solvent, such as an alkanol, ketone or ester. On the other hand, if the compound of this invention is desired in the free base form, it is isolated from a basic final wash step, according to the usual practice. A typical technique for preparing hydrochlorides is to dissolve the free base in a suitable solvent and dry the solution thoroughly, as over molecular sieves, before bubbling hydrogen chloride gas through it.

The dose of a compound of this invention to be administered to a human is rather widely variable and subject to the judgment of the attending physician. It should be noted that it may be necessary to adjust the dose of a compound when it is administered in the form of a salt, such as a laureate, the salt forming moiety of which has an appreciable molecular weight. The general range of effective administration rates of the compounds is from about 0.05 mg/day to about

100 mg/day. A preferred rate range is from about 0.25 mg/day to 25 mg/day. Of course, it is often practical to administer the daily dose of compound in portions, at various hours of the day. However, in any given case, the amount of compound administered will depend on such factors as the solubility of the active component,
5 the formulation used and the route of administration.

The route of administration of the compounds of this invention is not critical. The compounds are known to be absorbed from the alimentary tract, and so it is usually preferred to administer a compound orally for reasons of convenience. However, the compounds may equally effectively be administered percutaneously, or
10 as suppositories for absorption by the rectum, if desired in a given instance.

The compounds of this invention are usually administered as pharmaceutical compositions which are important and novel embodiments of the invention because of the presence of the compounds. All of the usual types of compositions may be used, including tablets, chewable tablets, capsules, solutions,
15 parenteral solutions, troches, suppositories and suspensions. Compositions are formulated to contain a daily dose, or a convenient fraction of daily dose, in a dosage unit, which may be a single tablet or capsule or convenient volume of a liquid.

Any of the compounds may be readily formulated as tablets, capsules and the like; it is preferable to prepare solutions from water-soluble salts, such as the
20 hydrochloride salt. In general, all of the compositions are prepared according to known methods in pharmaceutical chemistry. Capsules are prepared by mixing the compound with a suitable diluent and filling the proper amount of the mixture in capsules. The usual diluents include inert powdered substances such as starch of many different kinds, powdered cellulose, especially crystalline and microcrystalline
25 cellulose, sugars such as fructose, mannitol and sucrose, grain flours and similar edible powders.

Tablets are prepared by direct compression, by wet granulation, or by dry granulation. Their formulations usually incorporate diluents, binders, lubricants and disintegrators as well as the compound. Typical diluents include, for example,

various types of starch, lactose, mannitol, kaolin, calcium phosphate or sulfate, inorganic salts such as sodium chloride and powdered sugar. Powdered cellulose derivatives are also useful. Typical tablet binders are substances such as starch, gelatin and sugars such as lactose, fructose, glucose and the like. Natural and
5 synthetic gums are also convenient, including acacia, alginates, methylcellulose, polyvinylpyrrolidone and the like. Polyethylene glycol, ethylcellulose and waxes can also serve as binders.

A lubricant is typically necessary in a tablet formulation to prevent the tablet and punches from sticking in the die. The lubricant is chosen from such
10 slippery solids as talc, magnesium and calcium stearate, stearic acid and hydrogenated vegetable oils. Tablet disintegrators are substances which swell when-wetted to break up the tablet and release the compound. They include starches, clays, celluloses, algin and gums. More particularly, corn and potato starches, methylcellulose, agar, bentonite, wood cellulose, powdered natural sponge, cation-
15 exchange resins, alginic acid, guar gum, citrus pulp and carboxymethyl cellulose, for example, may be used as well as sodium lauryl sulfate. Tablets are often coated with sugar as a flavor and sealant, or with film-forming protecting agents to modify the dissolution properties of the tablet. The compounds may also be formulated as chewable tablets, by using large amounts of pleasant-tasting substances such as
20 mannitol in the formulation, as is now well - established in the art.

When it is desired to administer a compound as a suppository, the typical bases may be used. Cocoa butter is a traditional suppository base, which may be modified by addition of waxes to raise its melting point slightly. Water-miscible suppository bases comprising, particularly, polyethylene glycols of various molecular
25 weights are in wide use.

The effect of the compounds may be delayed or prolonged by proper formulation. For example, a slowly soluble pellet of the compound may be prepared and incorporated in a tablet or capsule, or as a slow-release implantable device. The technique may be improved by making pellets of several different dissolution rates

and filling capsules with a mixture of the pellets. Tablets or capsules may be coated with a film which resists dissolution for a predictable period of time. Even the parenteral preparations may be made long-acting, by dissolving or suspending the compound in oily or emulsified vehicles which allow it to disperse only slowly in the
5 serum.


The following Examples are presented by way of illustration, not limitation.

EXAMPLES

In summary, Examples 1-11 are directed to the synthesis of
10 representative compounds of this invention. Example 12 discloses a human bone cell co-culture system for assaying compounds for their ability to block production of IL-6. Example 13 discloses the activity of a representative compound of this invention as assayed in the human bone co-culture system of Example 12. Example 14-21 presents further experimental data evidencing activity of a representative compound
15 of this invention.

EXAMPLE 1

2-(4-HYDROXYBENZYLACETONE)-5-METHOXYPHENOL




To a mixture of 3-methoxyphenol (50 g, 0.40 mol), 4-
20 hydroxyphenylacetic acid (71 g, 0.46 mol) and ZnCl_2 (174 g, 1.28 mol) was added POCl_3 (100 ml, 1.6 mol). The mixture was stirred at 65°C for 2 hours, poured into ice water (2 L) and stirred until the ice melted. The clear supernatant was decanted and the residue was rinsed with water (1 L) and partitioned between EtOAc and

water. The organic layer was washed with brine, dried (MgSO_4), filtered and concentrated. The resulting oil was purified by chromatography (SiO_2 , 20% EtOAc/n-hexane) to provide 2-(4-hydroxybenzylacetone)-5-methoxyphenol (34.1 g, 33% yield) as a white solid; mp 137-140°C.

5

EXAMPLE 2


2-(4-TRIISOPROPYLSILOXYBENZYLACETONE)-5-METHOXYPHENOL



To a mixture of 2-(4-hydroxybenzylacetone)-5-methoxyphenol (10 g, 0.038 mole), NEt_3 (6 ml, 0.042 mole) in CH_2Cl_2 (50ml) was added
10 triisopropylsilylchloride (9 ml, 0.042 mole). The mixture was stirred for 22 hours, concentrated and the residue partitioned between EtOAc and H_2O . The organic layer was washed with NaOH (1N), HCl (1N) and brine. The organic layer was dried (MgSO_4), filtered and concentrated. The residue was treated with n-hexane to
15 provide 2-(4-triisopropylsilyloxybenzylacetone)-5-methoxyphenol (6.2 g, 38% yield) as a white solid; mp 66-68 °C.

EXAMPLE 3


3-PHENYL-4-(4-HYDROXYBENZYL)-7-METHOXYCOUMARIN



To a mixture of 2-(4-triisopropylsilyloxybenzylacetone)-5-methoxyphenol (4 g, 9.6 mmole), K_2CO_3 (4 g, 29 mmole) in CH_3CN (50 ml) was added phenyl acetylchloride (2.3 ml, 14 mmole). The mixture was stirred at reflux for 22 hrs, poured into H_2O ($0^\circ C$) (500 ml) and extracted with EtOAc (2x). The organic layer was dried ($MgSO_4$), filtered and concentrated. The residue was stirred with Et_2O and the resulting solid was filtered and recrystallized (EtOH) to give 3-phenyl-4-(4-hydroxybenzyl)-7-methoxycoumarin (0.88 g, 15% yield) as a white solid; mp $235-236^\circ C$.

EXAMPLE 4


10 3-PHENYL-4-[4-(2-{PIPERIN-1-YL})ETHOXY]-BENZYL-7-METHOXYCOUMARIN



15 A mixture of 3-phenyl-4-(4-hydroxybenzyl)-7-methoxycoumarin (0.50 g, 1.39 mmole), K_2CO_3 (0.58 g, 4.18 mmole), 2-chloroethylpiperidine hydrochloride (0.41 g, 2.22 mmole) and acetone (50 ml) was heated at reflux for 6 hours. The solvent was concentrated to a solid which was partitioned between EtOAc and H_2O . The organic layer was washed with NaOH (1H), brine, dried ($MgSO_4$), filtered and concentrated. The residue was stirred with HCl (20% in EtOAc) and the solid filtered to provide 3-phenyl-4-[4-(2-{piperin-1-yl})ethoxy]-benzyl-7-methoxycoumarin (0.57 g, 87% yield); mp $171-172^\circ C$.

EXAMPLE 5


3-PHENYL-4-[4-(2-{PIPERIDIN-1-YL})ETHOXY]-BENZYL-7-HYDROXYCOUMARIN



A mixture of 3-phenyl-4-[4-(2-{piperidin-1-yl})ethoxy]-benzyl-7-methoxycoumarin (0.10 g, 0.20 mmole), HOAc (glacial) (15 ml) and HBr (48%, 15ml) was refluxed for 48 hours. The mixture was partitioned between EtOAc (120 ml) and NaOH (1N, 120ml) and the aqueous layer was washed with EtOAc. The aqueous layer was then acidified (conc. HCl, pH 1-2) and filtered to provide 3-phenyl-4-[4-(2-{piperidin-1-yl})ethoxy]-benzyl-7-hydroxycoumarin (0.88 g, 99% yield); mp 160-161°C.

EXAMPLE 6


3-(4-FLUOROPHENYL)-4-[4-(1-METHYLPYRROLIDYL-3-OXY)]-BENZYL-7-HYDROXYCOUMARIN



A solution 3-(4-fluorophenyl)-4-[(4-hydroxyphenyl)methyl]-7-methoxy-2H-chromen-2-one (0.27 g, 0.72 mmole), in 3 mL of CH₂Cl₂ was treated


with 3-hydroxy-1-methylpiperidine (0.42g, 3.6mmol), triphenylphosphine (0.94 g, 3.6 mmol), and diethyl azodicarboxylate (0.65 g, 3.6 mmol). The reaction mixture was stirred for 8 hours at 25°C then concentrated under reduced pressure. The crude product was dissolved in 4 mL of a 1:1 solution of HBr (48%, aqueous) and glacial acetic acid. The resulting solution was warmed at 90°C for 12 hours. The reaction mixture was concentrated and the resulting residue was neutralized with 10 mL of saturated aqueous NaHCO₃. The aqueous mixture was extracted with CH₂Cl₂ (3 x 15 mL) and the combined organic layer was dried (MgSO₄) then concentrated under reduced pressure. The product (106 mg, 32%) was isolated following purification by flash chromatography (SiO₂, CH₂Cl₂/MeOH, 10:1).

Alternatively, 3-(4-fluorophenyl)-4-[(4-hydroxyphenyl)methyl]-7-methoxy-2H-chromen-2-one is reacted with one of the following enantiomers (a) or (b) in the presence of PPh₃ and diethyl azodicarboxylate (DEAD), followed by HBr/HOAc, to yield the corresponding enantiomeric product.



EXAMPLE 7

2-HYDROXY-4-METHOXYPHENYL 4-HYDROXYPHENYL




A solution of 3-methoxyphenol (2.5 g, 20.14 mmol), 4-hydroxybenzoic acid (3.2 g, 23.16 mmol), and ZnCl₂ (8.78 g, 64.44 mmol) in 15 mL of POCl₃ was warmed 65°C for 2 hours. The resulting reaction mixture was poured

onto ice water (100 mL). The aqueous layer was extracted with CH_2Cl_2 and the combined organic layer was dried (MgSO_4) then concentrated under reduced pressure. The crude product purified by flash chromatography to provide the title compound (3.93 g, 80%) as an off-white solid (LC/MS = 244 ($\text{M}+\text{H}^+$)).

5

EXAMPLE 8


3-(4-CHLOROPHENYL)-4-(4-HYDROXYPHENYL)-7-METHOXY-2H-CHROMEN-2-ONE 2



A suspension of 2-hydroxy-4-methoxyphenyl 4-hydroxyphenyl ketone
10 (2.6 grams, 10.65 mmol), 4-chlorophenylacetic acid (4.0 g, 23.43 mmol), K_2CO_3 (4.4
g, 31.95 mmol), carbonyldiimidazole (3.8 g, 23.43 mmol), and 4-
dimethylaminopyridine (0.1 g) in 20 mL of DMF was warmed at 90°C for 5 hours.
The reaction mixture was poured into 150 mL of H_2O and stirred for 30 minutes. The
precipitated product was collected by filtration and the purified desired product was
15 isolated following flash chromatography (2.1 g, 52%, LC/MS = 379 ($\text{M}+\text{H}^+$)).

EXAMPLE 9

3-(4-CHLOROPHENYL)-7-METHOXY-4-[4-(2-PIPERIDYLETHOXY)PHENYL]-2H-
CHROMEN-2-ONE




The title compound was prepared from 3-(4-chlorophenyl)-4-(4-hydroxyphenyl)-7-methoxy-2H-chromen-2-one as described in Example 4 (LC/MS = 492 (M+H⁺)).

10

EXAMPLE 10

3-(4-CHLOROPHENYL)-7-HYDROXY-4-[4-(2-PIPERIDYLETHOXY)PHENYL]-2H-
CHROMEN-2-ONE



The title compound was prepared from 3-(4-chlorophenyl)-7-methoxy-4-[4-(2-piperidylethoxy)phenyl]-2H-chromen-2-one as described in Example 5 (LC/MS = 476 (M+H⁺)).

EXAMPLE 11

5


ADDITIONAL REPRESENTATIVE COMPOUNDS
























By the procedures set forth herein, the compounds of Table 1 may be prepared.

Table 1

10

Representative Compounds



No.	R ₁	R ₂ [†]	R ₃	n	p	LC/MS (M+H ⁺)
1	Phenyl		H	2	1	456
2	4-fluorophenyl		H	2	1	474
3	4-chlorophenyl		H	2	1	490
4	4-bromophenyl		H	2	1	533,535
5	3-chlorophenyl		H	2	1	490
6	2-chlorophenyl		H	2	1	490
7	3-methylphenyl		H	2	1	470
8	2-methylphenyl		H	2	1	470
9	4-methylphenyl		H	2	1	470
10	4-hydroxyphenyl		H	2	1	473
11	3-bromopyridin-3-yl		H	2	1	535,537
12	3,4-dichlorophenyl		H	2	1	524,526
13	Thiophen-2-yl		H	2	1	462
14	4-trifluoromethylphenyl		H	2	1	524
15	4-chlorophenyl		H	2	2	504
16	4-chlorophenyl		H	2	0	476
17	Phenyl		H	2	1	458
18	4-fluorophenyl		H	2	1	474
19	Phenyl		SO ₂ N(CH ₃) ₂	2	1	563
20	Phenyl		CON(CH ₃) ₂	2	1	527
21	Phenyl		CO(phenyl)	2	1	560
22	Phenyl		COCH ₃	2	1	498
23	Phenyl		COOCH ₂ CH ₃	2	1	528

24	4-fluorophenyl			2	1	585
25	4-fluorophenyl		$\text{CH}_2\text{CH}_2\text{N}(\text{CH}_3)_2$	2	1	574
26	4-chlorophenyl		H	2	1	490
27	4-chlorophenyl		H	2	1	490
28	4-fluorophenyl		H	1	1	533
29	4-fluorophenyl		H	2	1	457
30	4-fluorophenyl		H	0	1	460
31	4-fluorophenyl		H	1	1	460
32	4-fluorophenyl		H	0	1	460
33	4-chlorophenyl		SO_2NH_2	2	1	570
34	2,4-difluorophenyl		H	2	1	492
35	2,4-dichlorophenyl		H	2	1	524
36	4-chlorophenyl		SO_2CH_3	2	1	568
37	4-fluorophenyl		H	3	1	448
38	Phenyl		H	2	0	430
39	4-chlorophenyl		CH_3	2	0	490
40	4-chlorophenyl		H	2	0	462
41	4-chlorophenyl		H	2	0	476
42	4-chlorophenyl		H	3	0	490
43	4-chlorophenyl		H	3	0	490

[†]At the 4-position of the phenyl ring unless otherwise noted

[‡]At the 3-position of the phenyl ring

EXAMPLE 12

HUMAN BONE CELL CO-CULTURE SYSTEM

The generation of osteoblastic cells from normal adult femoral
trabecular bone from patients without evidence of metabolic bone disease was carried
5 out as previously described (Kung Sutherland et al., *Osteoporosis International*
5:335-343, 1995; Wong et al., *J Bone Miner* 5:803-813, 1990). Bone fragments were
cleaned of adherent tissue and blood and cultured for 4 weeks in medium containing
Ham's F12 supplemented with 28 mM HEPES, pH 7.4, 10% FCS, 1.1 mM CaCl_2 ,
2 mM glutamine, and 1% antibiotic-antimycotic agent. Upon confluency, the cells
10 were harvested and immortalized using a retroviral vector (pLXSN) containing the
neomycin resistance gene and the human papillomaviral (HPV18) E6 and E7
oncogenes (*International Journal of Oncology* 6:167-174, 1995). The immortalized
osteoblastic cells were selected in medium containing G418 (600 $\mu\text{g/ml}$).

The human U937 monocytic cell line (cloned from the commercially
15 available cells by serial dilution) was maintained in RPMI 1640 medium containing
10% FCS and 1% antibiotic-antimycotic agent.

The co-culture system was established as follows. Human osteoblastic
cells were plated in 96-well dishes at a density of 5×10^3 cells/well. U937 cells in
RPMI medium (5×10^4 cells/well) were layered over the osteoblasts. The formation
20 of osteoclast-like cells was induced with 100 nM PMA. To determine the effects of
compounds or the neutralizing antibodies to IL-6 (Sigma) and GM-CSF
(Pharmingen) on the formation of osteoclast-like cells, the compounds or the
antibodies were added 30 minutes prior to the addition of PMA. The cultures were
stopped after 96 hrs and processed for histochemical or immunohistochemical
25 analyses.

EXAMPLE 13ACTIVITY OF REPRESENTATIVE COMPOUND
IN THE HUMAN BONE CO-CULTURE

The compound of Example 5 (*i.e.*, Compound No. 1 of Table 1) was
5 assayed in the human bone co-culture system of Example 12 and found to have an
IC₅₀ with respect to IL-6 and GM-CSF of 10 nM and 38 nM, respectively.

EXAMPLE 14ACTIVITY OF REPRESENTATIVE COMPOUND
ON IL-6 AND GM-CSF PRODUCTION IN HOB CELLS

10 Human osteoblasts (HOB) were plated in 96-well dishes at a density of
7 x 10³ cells/well in regular HOB medium (Ham's F12 supplemented with 28 mM
HEPES, pH 7.4, 10% FCS, 1.1 mM CaCl₂, 2 mM glutamine, and 1% antibiotic-
antimycotic agent). The following day, the cells were treated with Compound 1 or
vehicle (0.2% DMSO) for 30 minutes prior to the addition of the combination of
15 IL-1 β (1 ng/ml) and TNF- α (10 ng/ml). Cultures were continued for 18 to 24 hrs.
IL-6 and GM-CSF levels in culture medium were measured using commercially
available ELISA kits (Endogen, Cambridge, MA). The results of this experiment are
set forth in Figure 1A for IL-6 inhibition, and Figure 1B for GM-CSF inhibition.
These figures also present the comparative data for diethylstilbestrol (DES) and
20 Raloxifene.

EXAMPLE 15ACTIVITY OF REPRESENTATIVE COMPOUND
ON IL-6 PRODUCTION IN RA CELLS

Primary rheumatoid arthritic synoviocytes were plated in 96-well
5 dishes at a density of 8×10^3 cells/well in phenol-red free DMEM supplemented with
2 mM glutamine, 1% antibiotic-antimycotic agent and 10% charcoal-stripped FCS.
The following day, the cells were treated with Compound No. 1 or vehicle (0.2%
DMSO) for 30 minutes prior to the addition of the combination of IL-1 β (2 ng/ml)
and TNF- α (2 ng/ml). Cultures were continued for 18 to 24 hrs. IL-6 levels in
10 culture medium were measured using commercially available Endogen ELISAs as
disclosed above. The results of this experiment are presented in Figure 2, as well as
the comparative activity at DES and Raloxifene.

EXAMPLE 16ACTIVITY OF REPRESENTATIVE COMPOUND
ON PROLIFERATION OF BREAST CANCER CELLS

15 MCF-7 breast carcinoma cells were plated in 24-well dishes at a
density of 5×10^3 cells/well in phenol-red free DMEM:F-12 (1:1) medium containing
1% antibiotics, 0.05% β -mercaptoethanol, 0.01% ethanolamine, 0.42 ng/ml sodium
selenite and 5% charcoal-stripped FCS. Compound No. 1 and vehicle (0.2% DMSO)
20 were added the following day and refreshed with media change every 48 hours.
Cultures were stopped 9 days later and proliferation assayed using the Cyquant kit
(Molecular Probes, Eugene, OR). The results of this experiment are presented in
Figure 3, including comparative data for Tamoxifen and Raloxifene, as well as for the
combination of the same with Estradiol.

EXAMPLE 17ACTIVITY OF REPRESENTATIVE COMPOUND
ON PROLIFERATION OF PROSTATE CARCINOMA CELLS

DU-145 prostate carcinoma cells were plated in 96-well dishes at a
5 density of 2×10^3 cells/well in phenol-red free MEM Eagles medium containing
Earles' Balanced salts, 1% antibiotic-antimycotic agent, 2 mM glutamine, 0.1 mM
non-essential amino acids, 1 mM sodium pyruvate, and 10% charcoal-stripped FCS.
Compound No. 1 and vehicle (0.2% DMSO) were added the following day and
refreshed with medium change every 48 hrs. Cultures were stopped 5 days later and
10 proliferation assayed using the Cyquant kit as disclosed above. The results of this
experiment are presented in Figure 4, with comparative data for DES.

EXAMPLE 18

SOLID PHASE ER BINDING ASSAY

Binding to human Estrogen Receptor (ER- α and ER- β) was evaluated
15 in a solid-phase ^3H -estradiol competition assay, as adapted from McGuire, *Cancer*
Res. 38:4289-4291, 1978. Briefly, human recombinant ER- α (15 nM) or ER- β (50
nM) was immobilized to wells of 96-well microtiter plates. Binding of test
compound to ER was evaluated in competition with 30 nM ^3H -estradiol at room
temperature for 1 hour. The results of this experiment, as presented in Table 2,
20 represent the average from at least 3 different experiments. Reference compounds
were tested concurrently as an integral part of each assay to ensure the validity of the
results obtained.

Table 2
Solid Phase ER Binding

Receptor Assay	K _i (nM)			
	Compound No. 1 (HCl salt)	Tamoxifen citrate	Raloxifene HCl	Estradiol
ER- α	1.4	72	0.4	0.8
ER- β	7.3	173	13	2.5

5 Binding data for tamoxifen, raloxifene, and estradiol were found to correspond to published literature values for these compounds. Binding of Compound No. 1 (in the form of the HCl salt) to ER- α and ER- β was compared to binding of 17 β -estradiol, tamoxifen citrate and raloxifene-HCl. Compound No. 1 binds with high affinity (similar to estradiol) to ER- α and ER- β . Affinity for ER- β was found to be slightly lower than from ER- α , as is typical for most ER ligands

EXAMPLE 19

UTERINE LIABILITY

As discussed herein, compounds that function as Selective Estrogen Receptor Modulators or SERMs are desirable. In a preferred embodiment, SERMs do not exhibit uterine liability as assessed in recognized assays, such as the immature rat uterine bioassay (Robertson et al., *Biol. Reprod.* 54:183-196, 1996; Medlock et al., *Biol. Reprod.* 56:1239-1244, 1997; Martel et al., *Endocrinology* 139:2486-2492, 1998; Hyder et al., *Cancer Cells* 120:165-171, 1997), and the ovariectomized rat uterine bioassay (Sato et al., *FASEB J.* 10905-912, 1996; Ruenitz et al., *Bone* 23:537-542, 1998; Luo et al., *Endocrinology* 139:2645-2656, 1998; Ke et al., *Bone* 20:31-39, 1997). Both of these assays measure the effect of a particular compound on uterine wet and dry weight, and provide histological data.

Compound No. 1 was tested side-by-side with estrogen and Raloxifene-HCl in a 3-day immature rat uterine bioassay using subcutaneous (s.c.) administration (Medlock et al., *supra*). Estrogen caused a 4.5-fold increase in uterine weight, which is the reported literature value (Ashby et al., *Regul. Toxicol. Pharmacol.* 25:226-231, 1997; Medlock et al., *supra*). Raloxifene-HCl caused up to 50% increase in uterine weight at the 1mg/kg dose, which is also in agreement with published data (Ashby et al., *supra*). Compound No. 1 caused about a 30% increase in uterine weight, which was not statistically different from the vehicle-treated animals (i.e., PEG-400). Both Raloxifene-HCl and Compound No. 1 displayed a bell-shaped dose response, which is typically found with SERMs.


Compound No. 1 was also tested side-by-side with 17 β -estradiol and Raloxifene-HCl in a 28-day uterine bioassay in ovariectomized rats (Ke et al., *supra*). All three compounds were administered daily by s.c. injection. Estrogen prevented the uterine weight loss caused by ovariectomy by increasing the uterine weight by 550%. These results are in agreement with published data (Grese et al., *Proc. Natl. Acad. Sci. USA* 94:14105-14110, 1997; Ashby et al., *supra*; Ke et al., *supra*). Raloxifene-HCl caused about a 70% increase in uterine weight which has been previously reported (Grese et al., *supra*; Ashby et al., *supra*). Compound No. 1 caused about a 40% increase in uterine weight, which was statistically less than the Raloxifene-HCl-mediated weight increase. These results support the data from the immature rat uterine bioassay that Compound No. 1 does not exhibit significant uterine liability. Thus, this compound has significant potential for the prevention and/or treatment of cancer and osteoporosis, as well as cardiovascular diseases and neurodegenerative diseases such as Alzheimer's disease.

EXAMPLE 20

COMPARATIVE TESTING

Compound No. 1 was tested against a compound of Lednicer et al., *J. Med. Chem.* 8:725-726, 1965. More specifically, this reference is directed to coumarin-based compounds as purported estrogen antagonists. One of these compounds (*i.e.*, Compound 20 of Table IV, hereinafter "L-20") has the following structure (LC/MS (M+H)⁺ = 414):

Compound "L-20"



The above compound was tested against Compound No. 38 of the present invention in the ER binding assay of Example 18 and the human bone co-
15 culture system of Example 12 for IL-6 activity. The results of these experiments are presented in Table 3.

Table 3

	ER Binding Assay		IL-6 Activity
	ER- α (K_i , μ M)	ER- β (K_i , μ M)	(IC_{50} , μ M)
Compound No. 38	0.69	1.56	0.175
"L-20"	>10	>10	>1

EXAMPLE 21

5

ER-SELECTIVITY IN U2OS OSTEOSARCOMA CELLS

Human U2OS osteosarcoma cells (ATCC) were stably transfected with expression vectors for human full-length ER- α or ER- β , respectively, using standard molecular biology techniques. Stable subclones were generated that expressed high levels of ER- α or ER- β mRNA. The expression of ER- α and ER- β was confirmed using RNase protection analysis. The parental U2OS cells did not express any measurable amounts of either ER- α or ER- β .

Cells were plated into 96-well plates at a density of 8000 cells per well in phenol red-free media with charcoal-stripped fetal calf serum. Twenty-four hours later, cells were either treated with vehicle (0.2% DMSO) or test compound at the concentrations indicated (in 0.2% DMSO). Thirty minutes later cells were stimulated with 10 ng/ml IL-1 β and 10 ng/ml TNF- α . Twenty-four hours later the media supernatant was analyzed for cytokine production (IL-6) using commercially available ELISA kits following the manufacturer's instructions. Cytokine production in the presence of vehicle (0.2% DMSO) was set to 100%.

Results of this assay are set forth in Table 4, for a number of representative compounds of this invention, as well as for the prior art test compounds. In this table, activity is expressed as IC_{50} calculated by 50% inhibition relative to DMSO control (100%).

Table 4

Effect of Representative SERMs on Cytokines


Compound	U2OS IL-6 ER- α IC ₅₀ (nM)	U2OS IL-6 ER- β IC ₅₀ (nM)	U2OS IL-6 ER-Minus IC ₅₀ (nM)
"L-20"	>1000	>1000	>1000
17- β -Estradiol	0.03	0.016	>1000
Raloxifene Hydrochloride	3	10,000	>1000
4-Hydroxy-Tamoxifen	0.3-3	>1000	>1000
Compounds No. 16	325	51	>1000
Compound No. 38	>1000	165	>10,000

Referring to the above table, representative compounds of this
5 invention (e.g., Compound Nos. 16 and 38) showed specificity for ER- β over ER- α ,
while the prior art compounds (with the exception of 17- β -Estradiol) were selective
for ER- α over ER- β .

It will be appreciated that, although specific embodiments of the
10 invention have been described herein for purposes of illustration, various
modifications may be made without departing from the spirit and scope of the
invention. Accordingly, the invention is not limited except by the following claims.

CLAIMS

1. A compound having the structure:



or a stereoisomer, prodrug or pharmaceutically acceptable salt thereof,

wherein:


n is 0, 1, 2, 3 or 4;

p is 0;

R_1 is an unsubstituted or substituted C_{6-12} aryl, C_{7-12} arylalkyl, C_{3-12} heterocycle or C_{4-16} heterocyclealkyl;

R_2 is NR_aR_b wherein R_a and R_b are independently hydrogen, C_{1-6} alkyl, C_{6-12} aryl, or heterocycle, and wherein R_a and R_b are optionally substituted with up to three substituents independently selected from C_{1-6} alkyl, halogen, C_{1-6} alkoxy, hydroxy and carboxyl;

or R_2 is a heterocyclic ring of the following structure:



wherein

m_1 and m_2 are independently 0, 1 or 2, and both m_1 and m_2 are not 0,

A is CH₂, O, S or NH,

Z represents 0, 1, 2 or 3 heterocyclic ring substituents selected from halogen, C₁₋₈alkyl, C₆₋₁₂aryl, C₇₋₁₂arylalkyl, C₃₋₁₂heterocycle, or C₄₋₁₆heterocyclealkyl,

and wherein any hydrogen atom on the heterocyclic ring may, taken together with a hydrogen atom on an adjacent atom of the heterocyclic ring, form a double bond;

R₃ is hydrogen, R₄, C(=O)R₄, C(=O)OR₄, CONHR₄, CONR₄R₅, or SO₂NR₅R₅;

R₄ and R₅ are independently C₁₋₃alkyl, C₆₋₁₂aryl, C₇₋₁₂arylalkyl, or a five- or six-membered heterocycle containing up to two heteroatoms selected from O, NR₆ and S(O)_q, wherein each of the above groups are optionally substituted with one to three substituents independently selected from R₇ and q is 0, 1 or 2;

R₆ is hydrogen or C₁₋₄alkyl; and


R₇ is hydrogen, halogen, hydroxy, C₁₋₆alkyl, C₁₋₄alkoxy, C₁₋₄acyloxy, C₁₋₄thio, C₁₋₄alkylsulfinyl, C₁₋₄alkylsulfonyl, (hydroxy)C₁₋₄alkyl, C₆₋₁₂aryl, C₇₋₁₂arylalkyl, COOH, CN, CONHOR₈, SO₂NHR₈, NH₂, C₁₋₄alkylamino, C₁₋₄dialkylamino, NHSO₂R₈, NO₂, or a five- or six-membered heterocycle, where each occurrence of R₈ is independently C₁₋₆alkyl.

2. The compound of claim 1 wherein R₁ is an unsubstituted or substituted C₆₋₁₂aryl.
3. The compound of claim 2 wherein R₁ is an unsubstituted or substituted phenyl.
4. The compound of claim 3 wherein R₁ is unsubstituted phenyl.

5. The compound of claim 3 wherein R_1 is substituted phenyl.
6. The compound of claim 5 wherein R_1 is 4-halophenyl, 4-methylphenyl, 4-hydroxyphenyl, 4-trifluorophenyl, 3-halophenyl, 2-halophenyl, 2,4-dihalophenyl, 3,4-dihalophenyl, wherein halo is fluoro, chloro, bromo or iodo.
7. The compound of claim 1 wherein R_1 is an unsubstituted or substituted heteroaryl.
8. The compound of claim 7 wherein R_1 is substituted or unsubstituted pyridinyl or thiophenyl.
9. The compound of claim 1 wherein R_1 is an unsubstituted or substituted C_{7-12} arylalkyl.
10. The compound of claim 9 wherein R_1 is an unsubstituted or substituted benzyl.
11. The compound of claim 1 wherein R_1 is an unsubstituted or substituted C_{3-12} heterocycle or C_{4-16} heterocyclealkyl.
12. The compound of claim 1 wherein n is 2.
13. The compound of claim 1 wherein n is 0, 1, 3 or 4.
14. The compound of claim 1 wherein R_3 is hydrogen.
15. The compound of claim 1 wherein R_3 is $C(=O)(C_{1-3}alkyl)$ or $C(=O)(C_{6-12}aryl)$.

16. The compound of claim 1 wherein R_1 is $C(=O)O(C_{1-8}\text{alkyl})$, SO_2NH_2 or $CONH_2$.

17. The compound of claim 1 wherein R_2 is a heterocyclic ring of the following structure:



18. The compound of claim 17 wherein m_1 and m_2 are 1.

19. The compound of claim 18 wherein A is CH_2 .

20. The compound of claim 19 wherein R_2 is piperidin-1-yl.

21. The compound of claim 18 wherein A is O.

22. The compound of claim 21 wherein R_2 is morpholin-4-yl.

23. The compound of claim 17 wherein m_1 is 1 and m_2 is 0.


24. The compound of claim 23 wherein A is CH_2 and R_2 is imidazolidin-2-yl substituted with 0 or 1 Z substituents.

25. The compound of claim 17 wherein R_2 is imidazol-1-yl or imidazol-2-yl substituted with 0 or 1 Z substituents.


26. The compound of claim 1 wherein the $R_2-(CH_2)_n-O-$ moiety is attached at the 4-position of the phenyl ring.

27. The compound of claim 1 wherein the $R_2-(CH_2)_n-O-$ moiety is attached at the 3-position of the phenyl ring.

28. The compound of claim 1 having the structure:




29. The compound of claim 1 having the structure:




wherein X represents one or more phenyl substituents.


30. The compound of claim 28 having the structure:



31. The compound of claim 29 having the structure:



32. The compound of claim 31 having the structure:



wherein X is halogen.

33. A pharmaceutical composition comprising a compound of claim 1 in combination with a pharmaceutically acceptable carrier or diluent.

34. A method for inhibiting a cytokine in an animal in need thereof, comprising administering to the animal an effective amount of the composition of claim 33.

35. The method of claim 34 wherein the cytokine is IL-6.

36. The method of claim 34 wherein the cytokine is GM-CSF.

37. A method for modulating ER- β in a cell expressing ER- β , comprising contacting the cell with an effective amount of a compound of claim 1.
38. The method of claim 37 wherein the cell preferentially expresses ER- β over ER- α .
39. The method of claim 38 wherein the cell is of bone, bladder, uterus, ovary, prostate, testis, epididymis, gastrointestinal tract, kidney, breast, eye, heart, vessel wall, immune system, lung, pituitary, hippocampus or hypothalamus cell.
40. A method for modulating ER- β in tissue expressing ER- β , comprising contacting the tissue with an effective amount of a compound of claim 1.
41. The method of claim 40 wherein the tissue preferentially expresses ER- β over ER- α .
42. The method of claim 41 wherein the tissue is tissue of bone, bladder, uterus, ovary, prostate, testis, epididymis, gastrointestinal (GI) tract, kidney, breast, eye, heart, vessel wall, immune system, lung, pituitary, hippocampus or hypothalamus.
43. A method for treating an estrogen-related condition, comprising administering to an animal in need thereof an effective amount of a pharmaceutical composition of claim 33.
44. The method of claim 43 wherein the estrogen-related condition is breast cancer, osteoporosis, endometriosis, cardiovascular disease, hypercholesterolemia, prostatic hypertrophy, prostatic carcinomas, obesity, hot flashes, skin effects, mood swings, memory loss, prostate cancer, menopausal syndromes, type-II diabetes, Alzheimer's disease, urinary incontinence, GI tract conditions, spermatogenesis, vascular protection after injury, endometriosis, learning and memory, CNS effects, plasma lipid levels, acne, hirsutism, solid cancers, multiple myeloma, lymphoma, hairloss, cataracts, natural hormonal imbalances, or adverse reproductive effects associated with exposure to environmental chemicals.

45. A method for treating a bone-resorbing disease in an animal in need thereof, comprising administering to the animal an effective amount of the composition of claim 33.

46. The method of claim 39 wherein the bone-resorbing disease is osteoporosis.

47. The method of claim 39 wherein the bone-resorbing disease is metastatic bone cancer, osteolytic lesions with an orthopedic implant, Paget's disease, or bone loss associated with hyperparathyroidism.

48. A method for treating cancer associated with IL-6 in an animal in need thereof, comprising administering to the animal an effective amount of the composition of claim 35.

49. The method of claim 48 wherein the cancer is breast cancer, prostate cancer, colon cancer, endometrial cancer, multiple myeloma, renal cell carcinoma, or cervical carcinoma.


50. A method for treating arthritis in an animal in need thereof, comprising administering to the animal an effective amount of the composition of claim 33.

51. The method of claim 50 wherein the arthritis is rheumatoid arthritis.

52. A method for modulating gene expression in a cell expressing ER- β , comprising contacting the cell with an effective amount of a compound of claim 1.


53. The method of claim 52 wherein the cell is of bone, bladder, uterus, ovary, prostate, testis, epididymis, gastrointestinal tract, kidney, breast, eye, heart, vessel wall, immune system, lung, pituitary, hippocampus or hypothalamus.

54. The compound of claim 1, having the structure:




or a pharmaceutically acceptable salt thereof.

55. The compound of claim 1, having the structure:




or a pharmaceutically acceptable salt thereof.

56. The compound of claim 1, having the structure:




or a pharmaceutically acceptable salt thereof.

57. The compound of claim 1, having the structure:



or a pharmaceutically acceptable salt thereof.

58. The compound of claim 1, having the structure:



or a pharmaceutically acceptable salt thereof.

59. A pharmaceutical composition comprising a therapeutically effective amount of a compound of any one of claims 54-57 or 58 and a pharmaceutically acceptable carrier or diluent.

60. A method for inhibiting a cytokine in an animal in need thereof, comprising administering to the animal an effective amount of the composition of claim 59.

61. The method of claim 60 wherein the cytokine is IL-6.

62. The method of claim 60 wherein the cytokine is GM-CSF.

63. A method for modulating ER- β in a cell expressing ER- β , comprising contacting the cell with an effective amount of a compound of any one of claims 54-57 or 58.

64. The method of claim 63 wherein the cell preferentially expresses ER- β over ER- α .

65. The method of claim 64 wherein the cell is of bone, bladder, uterus, ovary, prostate, testis, epididymis, gastrointestinal tract, kidney, breast, eye, heart, vessel wall, immune system, lung, pituitary, hippocampus or hypothalamus cell.

66. A method for modulating ER- β in tissue expressing ER- β , comprising contacting the tissue with an effective amount of a compound of any one of claims 54-57 or 58.

67. The method of claim 66 wherein the tissue preferentially expresses ER- β over ER- α .

68. The method of claim 67 wherein the tissue is tissue of bone, bladder, uterus, ovary, prostate, testis, epididymis, gastrointestinal (GI) tract, kidney, breast, eye, heart, vessel wall, immune system, lung, pituitary, hippocampus or hypothalamus.

69. A method for treating an estrogen-related condition, comprising administering to an animal in need thereof an effective amount of a pharmaceutical composition of claim 59.

70. The method of claim 69 wherein the estrogen-related condition is breast cancer, osteoporosis, endometriosis, cardiovascular disease, hypercholesterolemia, prostatic hypertrophy, prostatic carcinomas, obesity, hot flashes, skin effects, mood swings, memory loss, prostate cancer, menopausal syndromes, type-II diabetes, Alzheimer's disease, urinary incontinence, GI tract conditions, spermatogenesis, vascular protection after injury, endometriosis, learning and memory, CNS effects, plasma lipid levels, acne, hirsutism, solid cancers, multiple myeloma, lymphoma, hairloss, cataracts, natural

hormonal imbalances, or adverse reproductive effects associated with exposure to environmental chemicals.

71. A method for treating a bone-resorbing disease in an animal in need thereof, comprising administering to the animal an effective amount of the composition of claim 59.

72. The method of claim 65 wherein the bone-resorbing disease is osteoporosis.

73. The method of claim 65 wherein the bone-resorbing disease is metastatic bone cancer, osteolytic lesions with an orthopedic implant, Paget's disease, or bone loss associated with hyperparathyroidism.

74. A method for treating cancer associated with IL-6 in an animal in need thereof, comprising administering to the animal an effective amount of the composition of claim 61.

75. The method of claim 74 wherein the cancer is breast cancer, prostate cancer, colon cancer, endometrial cancer, multiple myeloma, renal cell carcinoma, or cervical carcinoma.

76. A method for treating arthritis in an animal in need thereof, comprising administering to the animal an effective amount of the composition of claim 59.

77. The method of claim 76 wherein the arthritis is rheumatoid arthritis.

78. A method for modulating gene expression in a cell expressing ER- β , comprising contacting the cell with an effective amount of a compound of any one of claims 54-57 or 58.

79. The method of claim 78 wherein the cell is of bone, bladder, uterus, ovary, prostate, testis, epididymis, gastrointestinal tract, kidney, breast, eye, heart, vessel wall, immune system, lung, pituitary, hippocampus or hypothalamus.




Fig. 1A




Fig. 1B




Fig. 2





Fig. 3



Compound [μM]

Fig. 4